



## Well Child Schedule\*

<b>Age</b>	<b>Immunizations &amp; Screening Tests</b>
Newborn	Hepatitis B if not given in hospital
2 weeks	None
1 month	None
2 months	Pediarix (DTaP-IPV-Hep B), Hib, Prevnar (pneumococcal), rotavirus
4 months	Pediarix (DTaP-IPV-Hep B), Hib, Prevnar (pneumococcal), rotavirus
6 months	Pediarix (DTaP-IPV-Hep B), Hib, Prevnar (pneumococcal), rotavirus
9 months	Lead screening or test
12 months	MMR, Varivax (chickenpox), Hepatitis A, Hemoglobin test
15 months	DTaP, Hib, Prevnar (pneumococcal)
18 months	Hepatitis A, lead screening or test
24 months	None, if previous series completed
3 years	No vaccines if previous series completed, lead screening or test
4 years	May start KG series- MMR, Varivax, DTaP+IPV, lead screening or test
5 years	KG series (if not completed at 4 y/o)- MMR, Varivax, DTaP+IPV, lead screening or test
6-10 years**	Recommend yearly check-up but no vaccines needed, lead screening or test (6 years old)
11 years**	Menactra #1 (MCV, meningococcal), Tdap, Gardasil (HPV--series of 2 if completed prior to age 15 years)
12 years	Gardasil (HPV) if not completed
16 years	Menactra (MCV)#2, Bexsero #1 (MenB, meningococcal) [MenB #2 given 1-12 months later]
13-21 years**	Recommend yearly check-up, shots only if series not done.

We also recommend a **yearly Influenza vaccine** in the fall for all children age 6 months and older.

\*We follow the Well Child and immunization schedules recommended by the American Academy of Pediatrics (AAP) and the Centers for Disease Control (CDC).

\*\*A screening non-fasting lipid panel will be performed once between 9-11 years of age and again between 16-18 years of age. This is a recommendation of the AAP to screen for those at risk for early-onset heart disease.